

(Following Paper ID and Roll No. to be filled in your Answer Book)

PAPER ID : 1106 Roll No.

--	--	--	--	--	--	--	--	--	--	--

B. Tech.

(Semester-I) Theory Examination, 2012-13

ENGINEERING MATHEMATICS-I

Time : 3 Hours]

[Total Marks : 100

Note : Attempt questions from each Section as per instructions. The symbols have their usual meaning.

Section-A

Attempt *all* parts of this question. Each part carries 2 marks. $2 \times 10 = 20$

1. (a) If $y = x^2 \cdot \exp(2x)$, determine $(y_n)_0$.
- (b) Find the radius of curvature for the curve $s = \log(\tan \psi + \sec \psi) + \tan \psi \sec \psi$, where ψ is the angle which the tangent at any point to the curve makes with the x -axis.
- (c) If $u(x, y) = (\sqrt{x} + \sqrt{y})^5$, find the value of

$$\left(x^2 \frac{\partial^2 u}{\partial x^2} + 2xy \frac{\partial^2 u}{\partial x \partial y} + y^2 \frac{\partial^2 u}{\partial y^2} \right).$$

(d) The formula, $V = kr^4$, says that the volume V of fluid flowing through a small pipe or tube in a unit of time at a fixed pressure is a constant times the fourth power of the tube's radius r . How will a 10% increase in r affect V ?

(e) Use Beta function to evaluate :

$$\int_0^{\infty} \frac{x^8(1-x^6)}{(1+x)^{24}} dx$$

(f) Changing the order of integration in the double integral :

$$I = \int_0^8 \int_{\pi/4}^2 f(x, y) dx dy \text{ leads to}$$

$$I = \int_r^s \int_p^q f(x, y) dy dx \text{ say,}$$

What is p ?

(g) If $\vec{F} = \frac{\vec{r}}{r^3}$, find $\text{curl } \vec{F}$.

(h) Using Green's theorem, evaluate the integral :

$$\oint_C (xy dy - y^2 dx),$$

where C is the square cut from the first quadrant by the lines $x=1, y=1$.

(i) If $\alpha_1, \alpha_2, \alpha_3, \dots, \alpha_n$ are the characteristics roots of the n -square matrix A and k is a scalar, prove that the characteristic roots of $[A - kI]$ are $\alpha_1 - k, \alpha_2 - k, \alpha_3 - k, \dots, \alpha_n - k$.

(j) Explain the working rule to find the inverse of a matrix A by elementary row or column transformations.

Section-B

Attempt any *three* parts of this question. Each part carries 10 marks. $10 \times 3 = 30$

2. (a) Find the values of a and b such that the

expansion of $\log(1+x) - \frac{x(1+ax)}{(1+bx)}$ in ascending

powers of x begins with the term x^4 and hence find this term.

(b) Locate the stationary points of :

$$x^4 + y^4 - 2x^2 + 4xy - 2y^2$$

and determine their nature.

(c) Evaluate :

$$\int_R \int (x-y)^4 \cdot \exp(x+y) \cdot dx dy,$$

where R is the square in the xy -plane with vertices at $(1, 0)$, $(2, 1)$, $(1, 2)$ and $(0, 1)$.

- (d) Verify the Gauss divergence theorem for :

$$\vec{F} = (x^2 - yz)\hat{i} + (y^2 - xz)\hat{j} + (z^2 - xy)\hat{k}$$

taken over the rectangular parallelepiped

$$0 \leq x \leq a, 0 \leq y \leq b, 0 \leq z \leq c.$$

- (e) Diagonalize the following matrix A :

$$A = \begin{bmatrix} 3 & -1 & 1 \\ -1 & 5 & -1 \\ 1 & -1 & 3 \end{bmatrix}.$$

Section-C

Attempt *all* questions of this Section. Attempt any *two* parts from each question. Each question carries 10 marks. $10 \times 5 = 50$

3. (a) If $y = \sin[\log(x^2 + 2x + 1)]$, prove that :

$$(x+1)^2 y_{n+2} + (2n+1)(x+1)y_{n+1} + (n^2 + 4)y_n = 0.$$

- (b) Trace the curve $y = x(x^2 - 1)$.

- (c) Show that the radii of curvature of the

curve $y^2 = \frac{x^2(a+x)}{(a-x)}$ at the origin are $\pm a\sqrt{2}$.

1106

(4)

4. (a) The rate of flow Q of water per second

over the sharp-edged notch of length l , the height of the general level of the water above the bottom of the notch being h , is

$$\text{given by the formula } Q = c \left(1 - \frac{h}{5}\right) h^{3/2},$$

where c is a constant. Show that for small error δh in the measurement of h , the error δQ in Q is :

$$\frac{1}{2} c (3l - h) h^{1/2} \cdot \delta h.$$

- (b) Show that the envelope of the family of parabolas.

$$\left(\frac{x}{a}\right)^{1/2} + \left(\frac{y}{b}\right)^{1/2} = 1,$$

under the condition $ab = c^2$ (a , b and c are constants), is a hyperbola whose asymptotes coincides the axes.

- (c) Expand $f(x, y) = y^x$ about $(1, 1)$ up to second degree terms and hence evaluate

$$(1.02)^{1.03}.$$

1106

(5)

5. (a) Find the mass of the solid bounded by the ellipsoid $\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1$ and the coordinate planes where the density at any point $P(x, y, z)$ is $kxyz$, where k is a constant.

(b) Prove that :

$$\sqrt{m \left(m + \frac{1}{2} \right)} = \frac{\sqrt{\pi} \Gamma(2m)}{(2)^{2m-1}}$$

(c) Evaluate :

$$\int_0^\infty \int_0^x x \cdot \exp \left(-\frac{x^2}{y} \right) \cdot dx \cdot dy.$$

6. (a) A particle moves along a plane curve such that its linear velocity is perpendicular to the radius vector. Show that the path of the particle is a circle.

(b) Find the directional derivative of v^2 , where $\vec{v} = xy^2\hat{i} + zy^2\hat{j} + xz^2\hat{k}$ at the point $(2, 0, 3)$ in the direction of the outward normal to the sphere $x^2 + y^2 + z^2 = 14$ at the point $(3, 2, 1)$.

(c) Evaluate $\int_C \vec{F} \cdot d\vec{r}$ along the curve $x^2 + y^2 = 1$, $z = 1$ in the positive direction from $(0, 1, 1)$ to $(1, 0, 1)$, where :

$$\vec{F} = (yz + 2x)\hat{i} + xz\hat{j} + (xy + 2z)\hat{k}.$$

7. (a) Find the characteristic equation of the matrix :

$$A = \begin{bmatrix} 2 & 1 & -1 \\ 0 & 1 & 0 \\ 1 & 1 & 2 \end{bmatrix}$$

and hence find the matrix represented by

$$A^8 - 5A^7 + 7A^6 - 3A^5 + A^4 - 5A^3 + 8A^2 - 2A + I,$$

where $I = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$.

(b) Investigate for what values of λ, μ the simultaneous equations :

$x + y + z = 6$, $x + 2y + 3z = 10$, $x + 2y + \lambda z = \mu$ have (i) no solution, (ii) a unique solution (iii) an infinite number of solutions.

(c) If $N = \begin{bmatrix} 0 & 1+2i \\ -1+2i & 0 \end{bmatrix}$ is a matrix, then

show that $(I-N)(I+N)^{-1}$ is unitary

matrix, where $I = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$.